

### 3.2.2 死因

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=583	BU(経口)を含むレジメン n=199	
拒絶・生着不全	19 (3.3%)	6 (3.0%)	P=0.21
間質性肺炎以外の感染症	95 (16.3%)	36 (18.1%)	
間質性肺炎	39 (6.7%)	10 (5.0%)	
ARDS	7 (1.2%)	4 (2.0%)	
急性GVHD	30 (5.2%)	9 (4.5%)	
慢性GVHD	4 (0.7%)	7 (3.5%)	
VOD	16 (2.7%)	4 (2.0%)	
TMA	17 (2.9%)	8 (4.0%)	
原疾患の再発・増悪	249 (42.7%)	71 (35.7%)	
二次性悪性腫瘍	1 (0.2%)	1 (0.5%)	
出血	27 (4.6%)	9 (4.5%)	
外因死(自殺・事故含む)	1 (0.2%)	1 (0.5%)	
臓器不全(上記以外の原因による)	58 (9.9%)	29 (14.6%)	
その他	20 (3.4%)	4 (2.0%)	

	カテゴリ区分②				検定
	BU(経静脈) +CY±other drugs n=185	CY+TBI ±other drugs n=706	BU(経静脈) +FL±other drugs n=348	FL+TBI ±other drugs n=523	
拒絶・生着不全	6 (3.2%)	23 (3.3%)	11 (3.2%)	26 (5.0%)	P=0.04
間質性肺炎以外の感染症	31 (16.8%)	137 (19.4%)	53 (15.2%)	100 (19.1%)	
間質性肺炎	13 (7.0%)	45 (6.4%)	24 (6.9%)	28 (5.4%)	
ARDS	3 (1.6%)	15 (2.1%)	4 (1.2%)	12 (2.3%)	
急性GVHD	9 (4.9%)	26 (3.7%)	15 (4.3%)	30 (5.7%)	
慢性GVHD	3 (1.6%)	11 (1.6%)	1 (0.3%)	6 (1.2%)	
VOD	5 (2.7%)	29 (4.1%)	10 (2.9%)	8 (1.5%)	
TMA	5 (2.7%)	25 (3.5%)	10 (2.9%)	17 (3.3%)	
原疾患の再発・増悪	82 (44.3%)	287 (40.7%)	148 (42.5%)	167 (31.9%)	
二次性悪性腫瘍	0 (0.0%)	2 (0.3%)	1 (0.3%)	2 (0.4%)	
出血	7 (3.8%)	21 (3.0%)	18 (5.2%)	32 (6.1%)	
外因死(自殺・事故含む)	1 (0.5%)	3 (0.4%)	0 (0.0%)	0 (0.0%)	
臓器不全(上記以外の原因による)	13 (1.6%)	62 (8.8%)	41 (11.8%)	73 (14.0%)	
その他	7 (3.8%)	20 (2.8%)	12 (3.4%)	22 (4.2%)	