

3.4.2 性腺障害

男性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=743	BU(経口)を含むレジメン n=211	
性腺機能障害無	44 (5.9%)	15 (7.1%)	$P=0.36$
性腺機能障害有	5 (0.7%)	2 (0.9%)	
未検査	466 (62.7%)	118 (55.9%)	
不明	228 (30.7%)	76 (36.0%)	

	カテゴリ区分②				検定
	BU(経静脈) +CY±other drugs n=264	CY+TBI ±other drugs n=1116	BU(経静脈) +FL±other drugs n=436	FL+TBI ±other drugs n=677	
性腺機能障害無	22 (8.3%)	111 (10.0%)	18 (4.1%)	56 (8.3%)	$P=0.01$
性腺機能障害有	2 (0.8%)	7 (0.6%)	3 (0.7%)	1 (0.1%)	
未検査	173 (65.5%)	667 (59.8%)	266 (61.0%)	405 (59.8%)	
不明	67 (25.4%)	331 (29.7%)	149 (34.2%)	215 (31.8%)	

女性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=539	BU(経口)を含むレジメン n=161	
性腺機能障害無	44 (8.2%)	18 (11.2%)	$P=0.28$
性腺機能障害有	28 (5.2%)	8 (5.0%)	
未検査	298 (55.3%)	76 (47.2%)	
不明	169 (31.4%)	59 (36.6%)	

	カテゴリ区分②				検定
	BU(経静脈) +CY±other drugs n=201	CY+TBI ±other drugs n=811	BU(経静脈) +FL±other drugs n=287	FL+TBI ±other drugs n=408	
性腺機能障害無	12 (6.0%)	58 (7.2%)	27 (9.4%)	42 (10.3%)	$P < 0.001$
性腺機能障害有	10 (5.0%)	91 (11.2%)	14 (4.9%)	17 (4.2%)	
未検査	127 (63.2%)	418 (51.5%)	150 (52.3%)	214 (52.5%)	
不明	52 (25.9%)	244 (30.1%)	96 (33.4%)	135 (33.1%)	