

3.2.2 死因

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=1051	BU(経口)を含むレジメン n=228	
拒絶・生着不全	25 (2.4%)	8 (3.5%)	P=0.03
間質性肺炎以外の感染症	183 (17.4%)	42 (18.4%)	
間質性肺炎	62 (5.9%)	11 (4.8%)	
ARDS	15 (1.4%)	4 (1.8%)	
急性GVHD	54 (5.1%)	11 (4.8%)	
慢性GVHD	12 (1.1%)	10 (4.4%)	
VOD	34 (3.2%)	5 (2.2%)	
TMA	31 (3.0%)	8 (3.5%)	
原疾患の再発・増悪	441 (42.0%)	82 (36.0%)	
二次性悪性腫瘍	1 (0.1%)	1 (0.4%)	
出血	54 (5.1%)	8 (3.5%)	
外因死(自殺・事故含む)	1 (0.1%)	1 (0.4%)	
臓器不全(上記以外の原因による)	106 (10.1%)	33 (14.5%)	
その他	32 (3.0%)	4 (1.8%)	

	カテゴリ区分②				検定
	BU(経静脈) +CY±other drugs n=305	CY+TBI ±other drugs n=1115	BU(経静脈) +FL±other drugs n=682	FL+TBI ±other drugs n=860	
拒絶・生着不全	7 (2.3%)	34 (3.1%)	16 (2.4%)	35 (4.1%)	P<.0001
間質性肺炎以外の感染症	45 (14.8%)	201 (18.0%)	127 (18.6%)	189 (22.0%)	
間質性肺炎	19 (6.2%)	76 (6.8%)	40 (5.9%)	47 (5.5%)	
ARDS	4 (1.3%)	28 (2.5%)	10 (1.5%)	20 (2.3%)	
急性GVHD	16 (5.3%)	40 (3.6%)	31 (4.6%)	42 (4.9%)	
慢性GVHD	7 (2.3%)	22 (2.0%)	5 (0.7%)	13 (1.5%)	
VOD	15 (4.9%)	43 (3.9%)	18 (2.6%)	10 (1.2%)	
TMA	9 (3.0%)	34 (3.1%)	20 (2.9%)	25 (2.9%)	
原疾患の再発・増悪	131 (43.0%)	456 (40.9%)	281 (41.2%)	283 (32.9%)	
二次性悪性腫瘍	0 (0.0%)	2 (0.2%)	1 (0.2%)	3 (0.4%)	
出血	12 (3.9%)	36 (3.2%)	40 (5.9%)	48 (5.6%)	
外因死(自殺・事故含む)	1 (0.3%)	4 (0.4%)	0 (0.0%)	0 (0.0%)	
臓器不全(上記以外の原因による)	27 (8.9%)	104 (9.3%)	74 (10.9%)	110 (12.8%)	
その他	12 (3.9%)	35 (3.1%)	19 (2.8%)	35 (4.1%)	