

3.4.2 性腺障害

男性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=538	BU(経口)を含むレジメン n=117	
性腺機能障害無	54 (10.0%)	10 (8.6%)	P=0.72
性腺機能障害有	9 (1.7%)	3 (2.6%)	
未検査	335 (62.3%)	69 (59.0%)	
不明	140 (26.0%)	35 (29.9%)	

	カテゴリ区分②				検定
	BU(経静脈)+CY±other drugs n=185	CY+TBI±other drugs n=863	BU(経静脈)+FL±other drugs n=318	FL+TBI±other drugs n=440	
性腺機能障害無	21 (11.4%)	113 (13.1%)	30 (9.4%)	60 (13.6%)	P=0.20
性腺機能障害有	3 (1.6%)	11 (1.3%)	6 (1.9%)	2 (0.5%)	
未検査	117 (63.2%)	558 (64.7%)	194 (61.0%)	273 (62.1%)	
不明	44 (23.8%)	181 (21.0%)	88 (27.7%)	105 (23.9%)	

女性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=377	BU(経口)を含むレジメン n=90	
性腺機能障害無	38 (10.1%)	14 (15.6%)	P=0.52
性腺機能障害有	25 (6.6%)	5 (5.6%)	
未検査	210 (55.7%)	47 (52.2%)	
不明	104 (27.6%)	24 (26.7%)	

	カテゴリ区分②				検定
	BU(経静脈)+CY±other drugs n=146	CY+TBI±other drugs n=664	BU(経静脈)+FL±other drugs n=195	FL+TBI±other drugs n=279	
性腺機能障害無	10 (6.9%)	58 (8.7%)	21 (10.8%)	35 (12.5%)	P<0.001
性腺機能障害有	11 (7.5%)	124 (18.7%)	10 (5.1%)	26 (9.3%)	
未検査	87 (59.6%)	348 (52.4%)	109 (55.9%)	152 (54.5%)	
不明	38 (26.0%)	134 (20.2%)	55 (28.2%)	66 (23.7%)	