

3.2.2 死因

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=1616	BU(経口)を含むレジメン n=264	
拒絶・生着不全	44 (2.7%)	11 (4.2%)	P = 0.03
間質性肺炎以外の感染症	301 (18.6%)	43 (16.3%)	
間質性肺炎	89 (5.5%)	17 (6.4%)	
ARDS	19 (1.2%)	4 (1.5%)	
急性GVHD	77 (4.8%)	15 (5.7%)	
慢性GVHD	28 (1.7%)	13 (4.9%)	
VOD	48 (3.0%)	7 (2.7%)	
TMA	49 (3.0%)	9 (3.4%)	
原疾患の再発・増悪	652 (40.3%)	89 (33.7%)	
二次性悪性腫瘍	4 (0.2%)	1 (0.4%)	
出血	85 (5.3%)	9 (3.4%)	
外因死(自殺・事故含む)	1 (0.1%)	1 (0.4%)	
臓器不全(上記以外の原因による)	162 (10.0%)	37 (14.0%)	
その他	57 (3.5%)	8 (3.0%)	

	カテゴリ区分②				検定
	BU(経静脈) +CY±other drugs n=410	CY+TBI ±other drugs n=1568	BU(経静脈) +FL±other drugs n=1120	FL+TBI ±other drugs n=1254	
拒絶・生着不全	10 (2.4%)	42 (2.7%)	29 (2.6%)	59 (4.7%)	P < .0001
間質性肺炎以外の感染症	61 (14.9%)	279 (17.8%)	227 (20.3%)	268 (21.4%)	
間質性肺炎	20 (4.9%)	107 (6.8%)	65 (5.8%)	70 (5.6%)	
ARDS	5 (1.2%)	32 (2.0%)	13 (1.2%)	23 (1.8%)	
急性GVHD	19 (4.6%)	54 (3.4%)	51 (4.6%)	60 (4.8%)	
慢性GVHD	13 (3.2%)	34 (2.2%)	15 (1.3%)	20 (1.6%)	
VOD	18 (4.4%)	65 (4.1%)	28 (2.5%)	16 (1.3%)	
TMA	12 (2.9%)	44 (2.8%)	35 (3.1%)	34 (2.7%)	
原疾患の再発・増悪	181 (44.1%)	641 (40.9%)	431 (38.5%)	425 (33.9%)	
二次性悪性腫瘍	0 (0.0%)	3 (0.2%)	4 (0.4%)	8 (0.6%)	
出血	16 (3.9%)	52 (3.3%)	66 (5.9%)	60 (4.8%)	
外因死(自殺・事故含む)	1 (0.2%)	4 (0.3%)	0 (0.0%)	2 (0.2%)	
臓器不全(上記以外の原因による)	34 (8.3%)	143 (9.31)	121 (10.8%)	154 (12.3%)	
その他	20 (4.9%)	68 (4.3%)	35 (3.1%)	55 (4.4%)	