

3.4.2 性腺障害

男性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=825	BU(経口)を含むレジメン n=128	
性腺機能障害無	87 (10.6%)	11 (8.6%)	$P = 0.75$
性腺機能障害有	13 (1.6%)	3 (2.3%)	
未検査	505 (61.2%)	76 (59.4%)	
不明	220 (26.7%)	38 (29.7%)	

	カテゴリ区分②				検定
	BU(経静脈)+CY±other drugs n=279	CY+TBI±other drugs n=1184	BU(経静脈)+FL±other drugs n=497	FL+TBI±other drugs n=609	
性腺機能障害無	26 (9.3%)	151 (12.8%)	56 (11.3%)	83 (13.6%)	$P = 0.15$
性腺機能障害有	4 (1.4%)	16 (1.4%)	9 (1.8%)	3 (0.5%)	
未検査	178 (63.8%)	748 (63.2%)	298 (60.0%)	360 (59.1%)	
不明	71 (24.5%)	269 (22.7%)	134 (27.0%)	163 (26.8%)	

女性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=590	BU(経口)を含むレジメン n=99	
性腺機能障害無	61 (10.3%)	15 (15.2%)	$P = 0.51$
性腺機能障害有	37 (6.3%)	5 (5.1%)	
未検査	331 (56.1%)	51 (51.5%)	
不明	161 (27.3%)	28 (28.3%)	

	カテゴリ区分②				検定
	BU(経静脈)+CY±other drugs n=202	CY+TBI±other drugs n=947	BU(経静脈)+FL±other drugs n=347	FL+TBI±other drugs n=413	
性腺機能障害無	18 (8.9%)	89 (9.4%)	35 (10.1%)	59 (14.3%)	$P < 0.001$
性腺機能障害有	16 (7.9%)	200 (21.2%)	16 (4.6%)	35 (8.5%)	
未検査	114 (56.4%)	464 (49.0%)	202 (58.2%)	221 (53.5%)	
不明	54 (26.7%)	194 (20.5%)	94 (27.1%)	98 (23.7%)	