

3.2.2 死因

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=2247	BU(経口)を含むレジメン n=296	
拒絶・生着不全	60 (2.7%)	11 (3.7%)	P = 0.001
間質性肺炎以外の感染症	433 (19.3%)	46 (15.6%)	
間質性肺炎	121 (5.4%)	18 (6.1%)	
ARDS	32 (1.4%)	4 (1.4%)	
急性GVHD	102 (4.5%)	15 (5.1%)	
慢性GVHD	42 (1.9%)	17 (5.7%)	
VOD	67 (3.0%)	7 (2.4%)	
TMA	64 (2.9%)	9 (3.0%)	
原疾患の再発・増悪	864 (38.5%)	104 (35.1%)	
二次性悪性腫瘍	12 (0.5%)	1 (0.3%)	
出血	124 (5.5%)	9 (3.0%)	
外因死(自殺・事故含む)	2 (0.1%)	2 (0.7%)	
臓器不全(上記以外の原因による)	230 (10.2%)	43 (14.5%)	
その他	94 (4.2%)	10 (3.4%)	

	カテゴリ区分②				検定
	BU(経静脈) +CY±other drugs n=524	CY+TBI ±other drugs n=2009	BU(経静脈) +FL±other drugs n=1619	FL+TBI ±other drugs n=1674	
拒絶・生着不全	12 (2.3%)	50 (2.5%)	43 (2.7%)	74 (4.4%)	P < .0001
間質性肺炎以外の感染症	81 (15.5%)	356 (17.7%)	334 (20.6%)	364 (21.7%)	
間質性肺炎	25 (4.8%)	136 (6.8%)	91 (5.6%)	90 (5.4%)	
ARDS	8 (1.5%)	41 (2.0%)	23 (1.4%)	31 (1.9%)	
急性GVHD	25 (4.8%)	72 (3.6%)	70 (4.3%)	68 (4.1%)	
慢性GVHD	15 (2.9%)	49 (2.4%)	27 (1.7%)	27 (1.6%)	
VOD	20 (3.8%)	70 (3.5%)	44 (2.7%)	20 (1.2%)	
TMA	14 (2.7%)	52 (2.6%)	48 (3.0%)	45 (2.7%)	
原疾患の再発・増悪	215 (41.0%)	827 (41.2%)	601 (37.1%)	563 (33.6%)	
二次性悪性腫瘍	0 (0.0%)	6 (0.3%)	12 (0.7%)	10 (0.6%)	
出血	22 (4.2%)	69 (3.4%)	99 (6.1%)	81 (4.8%)	
外因死(自殺・事故含む)	1 (0.2%)	6 (0.3%)	1 (0.1%)	5 (0.3%)	
臓器不全(上記以外の原因による)	49 (9.4%)	186 (9.3%)	171 (10.6%)	211 (12.6%)	
その他	37 (7.1%)	89 (4.4%)	55 (3.4%)	85 (5.1%)	