

3.4.2 性腺障害

男性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=1151	BU(経口)を含むレジメン n=142	
性腺機能障害無	152 (13.2%)	15 (10.6%)	$P = 0.84$
性腺機能障害有	27 (2.4%)	3 (2.1%)	
未検査	616 (53.5%)	79 (55.6%)	
不明	356 (30.9%)	45 (31.7%)	

	カテゴリ区分②				検定
	BU(経静脈)+CY±other drugs n=371	CY+TBI±other drugs n=1573	BU(経静脈)+FL±other drugs n=718	FL+TBI±other drugs n=822	
性腺機能障害無	44 (11.9%)	188 (12.0%)	98 (13.7%)	133 (16.2%)	$P < 0.001$
性腺機能障害有	7 (1.9%)	28 (1.8%)	18 (2.5%)	7 (0.9%)	
未検査	204 (55.0%)	957 (60.8%)	377 (52.5%)	444 (54.0%)	
不明	116 (31.3%)	400 (25.4%)	225 (31.3%)	238 (29.0%)	

女性

	カテゴリ区分①		検定
	BU(経静脈)を含むレジメン n=832	BU(経口)を含むレジメン n=105	
性腺機能障害無	95 (11.4%)	17 (16.2%)	$P = 0.55$
性腺機能障害有	69 (8.3%)	8 (7.6%)	
未検査	423 (50.8%)	52 (49.5%)	
不明	245 (29.5%)	28 (26.7%)	

	カテゴリ区分②				検定
	BU(経静脈)+CY±other drugs n=258	CY+TBI±other drugs n=1235	BU(経静脈)+FL±other drugs n=522	FL+TBI±other drugs n=589	
性腺機能障害無	23 (8.9%)	125 (10.1%)	65 (12.5%)	90 (15.3%)	$P < 0.001$
性腺機能障害有	22 (8.5%)	266 (21.5%)	39 (7.5%)	50 (8.5%)	
未検査	134 (51.9%)	571 (46.2%)	266 (51.0%)	301 (51.1%)	
不明	79 (30.6%)	273 (22.1%)	152 (29.1%)	148 (25.1%)	